DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

MISCELLANEOUS INVESTIGATIONS SERIES
I-1310-C



CORRELATION OF MAP UNITS

Asterisks indicate position and adjacent numbers indicate age in millions of years of dated units



DESCRIPTION OF MAP UNITS

Qa ALLUVIUM (HOLOCENE)—Gravel, sand, and silt of channel, point bar, and floodplain deposits along the larger watercourses

DEPOSITS IN CLOSED BASINS (HOLOCENE)

Qab Fine-grained alluvium of poorly defined ephemeral stream systems in the axial parts of intermontane basins. Includes deposits in the southern part of San Simon Valley

Qpl Fine-grained playa and lacustrine sediments; locally includes beach sand and gravel

Qe EOLIAN DEPOSITS (HOLOCENE AND PLEISTOCENE)—Silt and sand in sheet deposits and irregular dunes

QTa FAN AND PEDIMENT ALLUVIUM (HOLOCENE TO PLIOCENE)—Grades into colluvial aprons along mountain fronts

QTaf Very coarse alluvium deposited in part as debris flows

QTar RIVER ALLUVIUM (PLEISTOCENE AND UPPER PLIOCENE)—Alluvium, typically silt, sand, and well-rounded gravel of widely dispersed provenance, forming the Gila River terrace system at elevations of ancestral Duck Creek (?) west and northwest of CHF 3LM

QTb ALKALI BASALT (PLEISTOCENE TO UPPER MIOCENE)—Generally olivine-bearing and pyroxene-bearing basalt; may contain some olivine, and pyroxene sanocrysts and mafic to ultramafic xenolithic nodules. Forms flows, plugs, and some cinder deposits

GILA CONGLOMERATE (PLEISTOCENE TO UPPER MIOCENE)—

QTg Fluvial and lacustrine deposits, generally locally derived and deposited in closed basins. Interbedded with **QTb**

QTgf Fine-grained alluvium, lacustrine deposits—Contains interculated lenses of coarser grained alluvium and thin beds of volcanic ash

QTgm Medium-grained alluvium, sandy and silty and fine gravel—Deposits in dated parts of alluvial fans and piedmonts. Includes lenses of coarser and fine grained alluvium

QTgc Coarse-grained alluvium and conglomerate—Deposits mainly in proximal parts of alluvial fans and piedmonts. Includes debris flow deposits and locally derived boulder conglomerate

Tvc VOLCANIC CONGLOMERATE (LOWER MIOCENE)—Chiefly well indurated, light-brown to light-gray, crudely bedded, matrix-supported conglomerate containing abundant volcanic clasts. Tuffaceous matrix commonly devitrified. Includes intercalated andesite flows and rhyolite ash-flow tuff beds. Deposited in basins in part different from those of today

Ta ANDESITE (MIOCENE AND OLIGOCENE)—Includes basalt, basaltic andesite, and dacite flows, and local intercalations of silicic pyroclastic deposits and volcaniclastic rocks; erupted from several largely unknown centers throughout the quadrangle. Forms extensive terranes. Most rocks are chemically porphyritic, and some of the dacitic units are strongly altered

Tr RHYOLITE (MIOCENE AND OLIGOCENE)—Forms dikes, plugs, and domes. Includes some intercalated andesite flows and massive flows. Chiefly flesh, fine-grained equigranular to porphyritic rhyolite and dacite. Some rocks are flow banded and contain spherulites and flow breccia

Ttg SACATON MOUNTAIN STOCK (MIOCENE)—

Tsc ROCKS OF THE SYCAMORE CANYON CALDERA (MIOCENE)—

Tsc Rhyolite flows, domes, and pyroclastic deposits with intercalated tuffaceous sandstone beds

Tsc PORPHYRITIC ANDESITE (MIOCENE AND OLIGOCENE)—Flows of coarse feldspar porphyry that resembles the turkey-track porphyry of Cooper (1960)

Tar ROCKS OF THE ASH PEAK-RHYOLITE PEAK ERUPTIVE CENTER (MIOCENE AND OLIGOCENE)—Chiefly rhyolite ash-flow tuff, domes, flows, and pyroclastic deposits and porphyritic dacite to rhyodacite domes and flows

Tgt ROCKS OF THE GREENWOOD MOUNTAIN ERUPTIVE CENTER (MIOCENE AND OLIGOCENE)

Tgr Rhyolite ash-flow tuff sheets, thin vitrophyric flows and interbedded volcaniclastic units

Tgd Porphyritic rhyolite domes and massive flows, and flow breccias

Tgd ROCKS OF THE BEAR SPRINGS VOLCANO (MIOCENE AND OLIGOCENE)—

Tgd Crystal-poor rhyolite dome

Tgr Rhyolite domes, flows, and dacite to rhyolite flows—Includes massive rhyolite flow and dacite flows at Bonita Creek

Tw TUFF OF WHATHERBY (MIOCENE AND OLIGOCENE)—

Tr Crystal-poor and crystal-rich ash-flow tuff and intercalated air-fall deposits and some rhyolite flows

Tpd RHYOLITE OF CANYON FORMATION (OLIGOCENE)—Chiefly rhyolite ash-flow tuff and intercalated volcaniclastic rocks and a rhyolite flow

Tpd ROCKS OF THE POWERLINE PASS AND DRY MOUNTAIN ERUPTIVE CENTERS (OLIGOCENE)—Chiefly andesite to rhyolite flows and pyroclastic deposits at Powerline Pass center and high-silica rhyolite dome-flow and pyroclastic deposits at Dry Mountain center

Tfe FELSIC VOLCANIC ROCKS (OLIGOCENE)—Mainly rhyolite, at scattered small localities

Tfi FELSIC TO INTERMEDIATE VOLCANIC ROCKS (OLIGOCENE)—Rhyolite to andesite flows, at scattered small localities

Tbc BLOODGOOD CANYON TUFF (OLIGOCENE)—Partly to densely welded, moderately crystal-rich, rhyolite ash-flow tuff derived from Bursum caldera north of Silver City quadrangle. In the Summit Mountains, may include older ash-flow tuff units from different sources

Tdr DACITE AND RHYOLITE (OLIGOCENE)—Chiefly thick, silicic flows, possibly related to rhyolite of unit Trh

Trh RHYOLITE (OLIGOCENE)—Large dome-flow complex near Tillie Hall Peak and smaller volcanic flows and intrusive rocks to the southeast

Tf FARAWAY RANCH FORMATION (OLIGOCENE)—Mainly rhyodacite flows and some intercalated andesite flows and pyroclastic and volcaniclastic rocks, and small intrusive masses

Tg GRANODIORITE OF GOLDEN GATE (OLIGOCENE)—Chiefly medium-grained, subhedral granular, biotite- or biotite-hornblende-bearing plutonic rocks and associated fine-grained porphyritic rocks. Forms dikes, sills, and small irregular intrusive masses. Include the Circle I Hills, Ninemile Canyon, and Buckhorn Basin plutons in the Dos Cabezas and Chiricahua Mountains and the Granite Gap pluton in Peloncillo Mountains

Tdb PYROCLASTIC AND VOLCANICLASTIC ROCKS (OLIGOCENE)—Crystal-rich ash-flow tuff, tuffaceous sandstone, and air-fall tuff. Includes the Caballo Blanco Rhyolite Tuff Member of Duff Formation and some rhyolite flows

Tdd DACITE (OLIGOCENE)—Massive biotite-hornblende dacite domes of Orange Butte and McKenzie Peak in the Peloncillo Mountains

Tw PORPHYRITIC ANDESITE (OLIGOCENE)—Fine- to medium-grained, locally vesicular, commonly porphyritic intrusive andesite

Td DACITE AND LATITE (OLIGOCENE)—Porphyritic flows with intercalated ash-flow tuff and tuffaceous sandstone. Unit may include flows of more widely diverse ages than indicated

Trh RHYOLITE (OLIGOCENE)—Massive, slightly porphyritic flow and subordinate sandstone at Steeple Rock in the Summit Mountains

Tc ROCKS OF THE CHIRICAHUA CALDERA COMPLEX (OLIGOCENE)—Mainly rhyolite flows, ash-flow tuff beds, some air-fall tuff beds, volcaniclastic units and andesite flows

Thm ROCKS OF THE HORSECAMP ERUPTIVE CENTER (OLIGOCENE)—Rhyolite dome, dacite breccia, and andesite to rhyolite flows. Includes massive rhyodacite flow at Midway Peak. May also include flows related to dome of Orange Butte

Tcr ROCKS OF STEINS VOLCANO (OLIGOCENE)

Tst Massive rhyolite and rhyodacite domes, flows, and breccias

Tst Rhyolite ash-flow tuff, partly filling a shallow subsidence structure

Tsf Massive rhyolite flows and air-fall tuff beds—Includes some intercalated andesite flows and massive dacite flows and breccia

Tdv PYROCLASTIC AND VOLCANICLASTIC ROCKS (OLIGOCENE)—Rhyolitic, crystal-rich ash-flow tuff, coarse sedimentary breccia, and air-fall tuff

Tn TADPOLE RIDGE FORMATION OF ELSTON AND OTHERS, 1973 (OLIGOCENE)—Densely welded rhyolitic ash-flow tuff

TBf ROCKS OF THE BESSIE RHOADES ERUPTIVE CENTER (OLIGOCENE)

Tbs Rhyolite flows and small domes and plugs

Tbs Rhyolite ash-flow and volcaniclastic rocks—May include some deposits from other sources

Tks KNEELING NUN AND SUGARLUMP TUFFS (OLIGOCENE)—Crystal-rich ash-flow tuff, Sugarlump Tuff present mainly in Silver City area; Kneeling Nun Tuff more widespread. Source probably the Emory caldera northeast of quadrangle

Tff LATITE AND PYROCLASTIC ROCKS (OLIGOCENE)—Chiefly latite flows and pyroclastic deposits and some intrusive rhyolite, ash-flow and air-fall deposits. Pyroclastic rocks are mainly crystal rich

Tr PYROCLASTIC AND VOLCANICLASTIC ROCKS (OLIGOCENE)—Mainly rhyolitic ash-flow tuff and sedimentary deposits, and some intercalated andesite and rhyolite flows. Source may be largely south of quadrangle

Tn NIPPER FORMATION OF SABINS (1957a) (OLIGOCENE)—Chiefly rhyolitic ash-flow tuff and some porphyritic and volcaniclastic rocks

Tcs ROCKS OF THE CLIFF EROSIVE CENTER (OLIGOCENE)

Tcs Porphyritic andesite flows

Tct Thin to massive, rhyolitic to quartz latitic ash-flow tuff beds, air-fall deposits, volcaniclastic rocks, and some silicic flows. May include some rocks from other sources

Trb ROCKS OF THE BURRO CIENAGA ERUPTIVE CENTER (OLIGOCENE)—Massive rhyolite flows, domes, and pyroclastic rocks

Tpr PYROCLASTIC AND VOLCANICLASTIC ROCKS (OLIGOCENE)—Mainly rhyolitic nonwelded ash-flow tuffs, air-fall tuffs, volcaniclastic rocks, and some intercalated basalt and rhyolite flows

Tdc DACITE (OLIGOCENE)—Plug of hornblende dacite porphyry containing sparse, large quartz phenocrysts

TRJbp BURRO PEAK FORMATION (OLIGOCENE)—Andesite to rhyodacite flows and breccia, conglomerate, and some tuffaceous sandstone

Tmp DIORITE (OLIGOCENE)—Fine-grained diorite plug

ROCKS OF THE MUIR ERUPTIVE CENTER (OLIGOCENE)

Tmc Precaldera rocks—Rhyolitic ash-flow tuff beds, ring-fracture latite and rhyolite domes and flows, and some andesite flows, volcaniclastic rocks, and lacustrine deposits

Tmc Precaldera and caldera rocks—Rhyolite flows and pyroclastic rocks from precaldera eruptions, and rhyolite ash-flow tuff and megabreccias from caldera-forming eruption

Tel ROCKS OF THE JPS MOUNTAIN ERUPTIVE CENTER (OLIGOCENE)—Andesite to dacite domes, flows, and volcaniclastic rocks

Tgg GRANITE (OLIGOCENE)—Medium-grained, porphyritic biotite granite of Gillespie Mountain pluton in the Peloncillo Mountains

Trhy RHYOLITE (OLIGOCENE AND EOCENE)—Mainly dikes of sparsely porphyritic rhyolite and other felsic rocks, chiefly in the Big Burro Mountains. Strongly altered. May include some older rocks in the Dragoon and Pyramid Mountains

Tad ANDESITE AND DACITE (OLIGOCENE AND EOCENE)—Mainly flows and some intercalated altered tuff, agglomerate, and volcaniclastic rocks, derived from dispersed centers, such as the Antarney Mountain eruptive center. Includes some rocks of Paleocene age in the Dragoon Mountains. Unit also includes irregular andesitic intrusive masses north of Silver City in the Pinos Altos Range

Tqs QUARTZ MONZONITE (EOCENE)—Medium grained, porphyritic, hypidiomorphic-granular amphibole quartz monzonite; locally biotite and pyroxene bearing. Includes the Silver City, Gomez Peak, and some smaller plutons along the southwest flank of the Pinos Alto Range

TKa ANDESITE (PALEOCENE AND UPPER CRETACEOUS)—Aphanitic and porphyritic flows and breccias, and some porphyritic and volcaniclastic rocks. Includes some intrusive andesite and dacite, chiefly sills and dikes, in scattered centers north to the Dos Cabezas volcanic field

TKg GRANODIORITE AND QUARTZ MONZONITE (PALEOCENE AND UPPER CRETACEOUS)—Medium-grained biotite granodiorite to plutons, plugs, and dikes. Includes the Silver Camp pluton (Dos Cabezas Mountains), Lone Star pluton (Peloncillo Mountains), Lordsburg pluton (Pyramid Mountains), Tyrone pluton (Big Burro Mountains), and the Pinos Altos and Pierce-Hanover plutons (Pinos Altos Range)

TKgd INTERMEDIATE INTRUSIVE ROCKS (PALEOCENE AND UPPER CRETACEOUS)—Mainly porphyritic dacite, quartz-diorite, monzonite, and syenodiorite in sills, laccoliths, dikes, and small plutons, chiefly in the Pinos Alto Range. Includes the Kneeling Nun pluton in the Dos Cabezas Mountains

Ksv SEDIMENTARY AND VOLCANIC ROCKS (UPPER CRETACEOUS)—Commonly includes andesite and volcaniclastic conglomerate with intercalated andesite flows and rhyolitic welded tuff. Conglomerate matrix coarse and subangular. Medium-assemble fossils of unit TKa. Welded tuff strongly altered, moderately indurated, contains chlorotized biotite

Kcb COLORADO FORMATION AND BEARTOOTH QUARTZITE (UPPER CRETACEOUS)—Dark-gray marine shale and sandstone and reddish-brown to white quartzite and quartz sandstone

Kb BISBEE GROUP (LOWER CRETACEOUS)—Dark-gray to pale-yellowish-brown, thin- to thick-bedded, fossiliferous limestone, sandstone, shale, limestone, and conglomerate. Commonly has a basal conglomerate unit. Includes marine, estuarine, and fluvial deposits

J'Jvs VOLCANIC AND SEDIMENTARY ROCKS (JURASSIC AND TRIASSIC)—Tuffaceous rocks, red shale, and conglomerate. Unit is volcaniclastic. Includes the Walnut Gap Formation

PPs SEDIMENTARY ROCKS (PERMIAN AND PENNSYLVANIAN)—Mainly limestone and some shale and sandstone. To the east includes Permian Concha Limestone, Scherrer Formation, Epitaph Dolomite, and Colina Limestone, and Pennsylvanian Horquilla Limestone and Naco Group. To the west includes Permian Epitaph Dolomite and Colina Limestone, and Pennsylvanian Syrena and Oswaldo Formations

MDs SEDIMENTARY ROCKS (MISSISSIPPIAN AND DEVONIAN)—Mainly limestone and some shale. Includes Mississippian Paradise Formation and Escabrosa Limestone, and Devonian Percha Shale to the east. Includes Devonian Percha Shale to the west

SOCs SEDIMENTARY ROCKS (SILURIAN TO CAMBRIAN)—Mainly dolomite, limestone, and sandstone, and some shale and limestone. Includes Silurian Fusselman Dolomite, Ordovician Montoya Group and El Paso Limestone, and Cambrian Bliss Sandstone and Lanoria Quartzite

OCs SEDIMENTARY ROCKS (ORDOVICIAN AND CAMBRIAN)—Dolomite, limestone, sandstone, and some shale, quartzite, and conglomerate. Includes Ordovician El Paso Limestone and Cambrian Concepcion Sandstone

Cs SEDIMENTARY ROCKS (CAMBRIAN)—Limestone, sandstone, shale, quartzite, and some dolomite and conglomerate. Includes Abrigo Formation and Bolsa Quartzite, and in the Dos Cabezas and Chiricahua Mountains includes a unit transitional between the El Paso and Abrigo

YXdp DIABASE (PROTEROZOIC Y)—Includes hornblende and some metadiorite. Forms dikes, plugs, and small plutons. Probably correlative with diabase of east-central Arizona dated at 1,200 m.y. (Silver, 1963)

Yg GRANODIORITE TO GRANITE (PROTEROZOIC Y)—Includes some aplite, pegmatite, and lamprophyre. Forms large plutons and dikes. Includes rocks of Rattlesnake Point (Chiricahua Mountains) and Big Burro Mountains plutons

YGgn GNEISS (PROTEROZOIC Y)—Hornblende and biotite gneiss, amphibolite, and migmatite. Probably largely of sedimentary origin. Includes Bullard Peak Series of Hewett (1959) in the Big Burro Mountains

YXgn GNEISS (PROTEROZOIC Y OR X)—Quartzofeldspathic gneiss that contains biotite or biotite and hornblende, and some amphibolite and schist. Probably of sedimentary and volcanic origin

Xm METASEDIMENTARY AND METAVOLCANIC ROCKS (PROTEROZOIC X)—Phyllite, schist, quartzite, and amphibolite; probably metabasite to meta-andesite, and some conglomerate. Locally includes some rocks similar to those in unit Ygn. Includes Pinal Schist of western area

CONTACT

FAULT—Dashed where inferred, dotted where concealed, queried where uncertain. Fault movements are indicated by the following symbols used in combination, as appropriate, for dome movement:

Normal fault—Ball and bar on downthrown side. In structure cross section arrow-coupled shows relative movement

Strike-slip fault—Arrow-couple shows relative movement. In structure cross section: A, movement away from viewer; T, movement toward viewer

Thrust fault—Sawteeth on upper plate

Subsurface fault—Inferred from geophysical data (Klein, unpub. data). M, aeromagnetic; G, gravity

STRANDLINE—Beach ridge, spit, and wave-cut benches of pluvial lakes, locally modified by alluvial or eolian processes. Hachures on lake side

VOLCANIC VENT—Chiefly local vent sites marked by pyroclastic cones in larger eruptive centers and some stratovolcanos and shield volcanoes. Additional vents may be inferred at some small intrusive bodies

GEOPHYSICAL DATA ON BASINS—Depth estimates are relative to other local basin features. Data from Klein (unpub. data)

DB Deep basin

B Basin

P Platform

1167† WELL LOCATION—Showing depth in feet and bedrock encountered. Bedrock symbols: Tv, Tertiary volcanic rocks; Tvc, Tertiary volcanic conglomerate; Kb, Cretaceous rocks; Pz, pre-Cretaceous rocks; Ig, granite; Ybd, Precambrian Y quartz monzonite; Xs, pre-Cretaceous rocks in Basin-fill at depth; ?, unknown



GEOLOGIC MAP OF THE SILVER CITY 1° x 2° QUADRANGLE, NEW MEXICO AND ARIZONA

By

Harald Drewes, B.B. Houser, D.C. Hedlund, D.H. Richter, C.H. Thorman, and T.L. Finnell

1985



INDEX MAP SHOWING SOME MAJOR TECTONIC AND PHYSIOGRAPHIC FEATURES IN ARIZONA, NEW MEXICO, AND MEXICO

For sale by Branch of Distribution, U.S. Geological Survey, Box